Last name _	
First name	

LARSON—MATH 353–CLASSROOM WORKSHEET 17 $n^2 + 1$ Primes Investigation.

- 1. Sign in to your CoCalc account.
 - (a) Start the Chrome browser.
 - (b) Go to https://cocalc.com
 - (c) Log in to your account.
 - (d) You should see an existing Project for our class. Click on that.
 - (e) Make sure you are in your Home directory (if you put files in the Handouts directory they could be overwritten.)
 - (f) Click "New", then "Jupyter Notebook", then call it 353-c17.
 - (g) Make sure you have SAGE as the kernel.
 - (h) Look in your Home directory. You should see a conjecturing.py file and an expressions file AND today's Jupyter notebook.

Review

- 2. We need conjecturing.py loaded for our investigation. I added the command to the number_theory.sage file. Every time that file is loaded, the Conjecturing program will get loaded too.
- 3. Open Conjectures.

```
count_prime_divisors(x) <= digits10(x)
count_prime_divisors(x) <= 1/2*count_divisors(x)
count_prime_divisors(x) <= count_divisors(x) - 1
count_prime_divisors(x) >= digits10(x) - 1
count_prime_divisors(x) >= (1/digits10(x))
```

Are any of these resolved?

Experiments

The **question** is: are the infinitely many primes of the form $n^2 + 1$?

4. To use the Conjecturing program, we'll need some *invariants*. The ones we've coded in class are in the *number_theory.sage* file in your Handouts folder. We don't want to keep re-coding those. We can use this file as a permanent record of everything we've coded for this research. Copy or move this file to your Home directory.

5. We started with a few prime and non-prime integers of the form $n^2 + 1$.

- 6. How should we proceed? Here are 2 new ideas: (1) we can add theorems using the (optional) theory parameter for the conjecture procedure; and (2) we can add new invariants.
- 7. If it is true, for instance, that for any $n=x^2+1$ integer that count_prime_divisors (x) <= $1/2 \times \text{count_divisors}$ (x) is true, we can define that upper bound using a procedure-name, and then adding that theorem to a list of theorems that we can input to the program:


```
def half_count_divisors(x):
    return (1/2)*count_divisors(x)

theorems = [half_count_divisors]

conjs = conjecture(objects, invariants, invariant_of_interest, upperBound = True, theory = theorems, debug = True)
```

8. Here's an idea for new invariants. For any $n=x^2+1$ integer, we can also grab onto the x and define any invariants for *that* integer. These are also invariants for x. There are no wrong invariants—and also you can't tell ahead of time what invariants will show up in useful conjectures.

Let's define these, add them to our invariants list, and then re-run conjecture.



Getting your classwork recorded

When you are done, before you leave class...

- 1. Click the "Print" menu choice (under "File") and make a pdf of this worksheet (html is OK too).
- 2. Send me an email (clarson@vcu.edu) with an informative header like "Math 353 c17 worksheet attached" (so that it will be properly recorded).
- 3. Remember to attach today's classroom worksheet!